

Patient Name First Name \_\_\_\_\_ Middle Name \_\_\_\_\_ Last Name \_\_\_\_\_ DOB \_\_\_\_\_ Weight \_\_\_\_\_  Male  Female

Street Address \_\_\_\_\_ Apt # \_\_\_\_\_ City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Daytime Tel \_\_\_\_\_ Evening Tel \_\_\_\_\_ Cell \_\_\_\_\_ Email \_\_\_\_\_

Ship to Patient at  Home  Work **OR** Patient will pick up at  Physician Office  Pharmacy Date Needed \_\_\_\_\_DIAGNOSIS  Polymyositis (M33.20)  CVID (D84.9)  CIDP (G61.81)  Other specified diabetes mellitus with diabetic neuropathy, unspecified (E13.40) Guillian-Barre Syndrome (G61.0)  Immune Neuropathy other than CIDP w/o Paraproteinemia (G61.81)  Dermatomyositis (M36.0)  Myathesnia Gravis (G70.0) Lambert-Eaton Syndrome, unspecified (G70.80)  Polyneuropathy in diseases classified elsewhere (G63)  Other \_\_\_\_\_Insured's Name \_\_\_\_\_ Relation to Patient \_\_\_\_\_ Eligible for Medicare  Yes  No If yes, Medicare# \_\_\_\_\_Prescription Card  Yes  No If Yes, Carrier \_\_\_\_\_ Tel \_\_\_\_\_ Fax \_\_\_\_\_ Policy/Group# \_\_\_\_\_

Bin# \_\_\_\_\_ Pcn# \_\_\_\_\_ RXID# \_\_\_\_\_ RX Group# \_\_\_\_\_

Prescriber's Name \_\_\_\_\_ Office Contact \_\_\_\_\_

Street Address \_\_\_\_\_ Suite # \_\_\_\_\_ City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Tel \_\_\_\_\_ Fax \_\_\_\_\_ Email \_\_\_\_\_

License# \_\_\_\_\_ NPI# \_\_\_\_\_ UPIN# \_\_\_\_\_ DEA# \_\_\_\_\_

### PRESCRIPTION

### PLEASE ATTACH COPIES OF PATIENT'S INSURANCE CARDS

Does patient already have a line?  Yes  No If yes, type of line \_\_\_\_\_IVIG to be infused via existing line:  Yes  No**First IVIG Infusion?** Yes If yes, IgA level is more than 5 mg/dl:  Yes  No  Not Available**Ig Quantification:** IgA, IgG, IgM (prior to 1<sup>st</sup> IVIG infusion) No If no, brand/dose of IVIG: \_\_\_\_\_ Last infusion Date: \_\_\_\_\_

Note: IVIG contains IgA and is contraindicated in IgA deficient patients with antibodies against IgA and history of hypersensitivity.

**IVIG (IMMUNOGLOBIN) ORDER:** \_\_\_\_\_ (IVIG brand will be chosen if not specified)**IVIG DOSE:** \_\_\_\_\_ g/kg = \_\_\_\_\_ g (round to nearest vial size) infuse intravenously (Range: 0.2-2 g/kg) Repeat dose **daily** x \_\_\_\_\_ consecutive days total, repeat dose: monthly x \_\_\_\_\_ months Repeat dose **weekly** x \_\_\_\_\_ weeks total Repeat dose **monthly** x \_\_\_\_\_ months total Other \_\_\_\_\_**SUPPLIES FOR INFUSION** (If Necessary) NaCl 0.9% / D5W for flush: flush Line/Port with (3 - 5 ml for PIV and 5-10 ml for Central Line/Port) per nursing agency protocol (NaCl 0.9% / D5W will be used based on IVIG compatibility) Heparin for flush (100 Units / ml) (if RN keeps PIV or if needed for Central Line), flush with 3-5 ml per nursing agency protocol Sterile water for reconstitution of powder to make the requested concentration (for Carimune NF) Other: \_\_\_\_\_**SUGGESTED RATE OF INFUSION:** 30 -150 ml/hr as tolerated (Increase rate gradually every 30 min by 20-30 ml/hr) Other \_\_\_\_\_**PRE-MEDICATIONS: TO BE ADMINISTERED 30 MIN PRIOR TO IVIG INFUSION (QTY:PER INFUSION):** Diphenhydramine 25 - 50 mg PO Dispense:#2 (25 mg) Acetaminophen 650 mg PO Dispense:#2 (325 mg) Other \_\_\_\_\_ QTY: QS**IN THE EVENT OF ANAPHYLAXIS:**

• Stop Infusion and call MD &amp; 911

• Diphenhydramine 25 - 50 mg IVP every 4 hours prn

(Not to exceed 25 mg/min) QTY: 3 (50 mg)

• Epinephrine (1:1000) 0.4 mg SQ prn anaphylaxis,

may repeat every 20 minutes x 2 QTY: 3 amp

Other \_\_\_\_\_

 **ENROLL IN NURSE TRAINING / MANUFACTURER PROGRAM****Prescriber's Signature** (signature required. NO STAMPS) \_\_\_\_\_ **Date** \_\_\_\_\_

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately. PLEASE NOTE: Bergen can only accept original prescription drug orders from patients, faxed prescriptions can be accepted only from the prescribing practitioners.